

SPECIAL LAGRANGIAN CONES IN $\mathbb{C}^3$ AND PRIMITIVE HARMONIC MAPS.

IAN MCINTOSH

ABSTRACT. In this article I show that every special Lagrangian cone in $\mathbb{C}^3$ determines, and is determined by, a primitive harmonic surface in the 6-symmetric space SU_3/SO_2 . For cones over tori, this allows us to use the classification theory of harmonic tori to describe the construction of all the corresponding special Lagrangian cones. A parameter count is given for the space of these, and some of the examples found recently by Joyce are put into this context.

1. INTRODUCTION.

Special Lagrangian submanifolds are the subject of a great deal of attention at present, largely due to their central role in a number of conjectures concerning the mirror symmetry of Calabi-Yau 3-folds (see, for example, [10, 14, 22]). In particular, some effort has been put into the construction of examples of special Lagrangian cones in $\mathbb{C}^3$ [11, 13], since these provide local models for the singularities which may occur in special Lagrangian fibrations of a Calabi-Yau 3-fold. It is not hard to see that a special Lagrangian cone in $\mathbb{C}^3$ determines, as its link (its intersection with the unit 5-sphere), a minimal Legendrian surface in S^5 and by Hopf projection a minimal Lagrangian surface in $\mathbb{CP}^2$, and (locally, at least) the reverse is true. Now, minimal surfaces in spheres and complex projective spaces possess a well-developed Riemannian twistor theory [4] which asserts that each may be lifted to a harmonic map into a flag manifold which possesses a sort of horizontal holomorphicity: this lift is called a primitive harmonic lift. The first aim of this note is to describe the relevant twistor theory for special Lagrangian cones in $\mathbb{C}^3$.

The second aim of this note is to describe the construction of all minimal Lagrangian tori in $\mathbb{CP}^2$, which is equivalent to the construction of special Lagrangian cones over tori. Here I adapt the construction of non-isotropic minimal tori given in [16, 17] to encode the extra condition of being Lagrangian. The construction of [16, 17] assigns to each minimal torus its spectral data $(X, \lambda, \mathcal{L})$ consisting of a real algebraic curve X , a degree three function λ on X and a line bundle $\mathcal{L}$ over X , with $\lambda, \mathcal{L}$ both respecting the real involution carried by X . The extra condition of being Lagrangian requires that X possess an extra holomorphic involution μ which $\lambda, \mathcal{L}$ must also respect (this much has also been noted by [21, 15]). In particular, $\mathcal{L}$ lives on a translate of the Prym variety $\text{Prym}(X, \mu)$.

1991 *Mathematics Subject Classification.* 53C43, 58E20.

My interest in this topic grew out of conversations with Dominic Joyce concerning his construction of special Lagrangian cones over tori [13]. As I became more involved it emerged that certain aspects of the problem had been tackled, from one angle or another, at least five times in the last decade. The unifying object is the p.d.e

$$(1) \quad \frac{\partial^2 u}{\partial z \partial \bar{z}} = e^{-2u} - e^u,$$

which is known either as Tzitzéica's equation or the Bullough-Dodd-Zhiber-Shabat equation. In our context it occurs as the equation for the metric $e^u |dz|^2$ induced on a torus by its minimal Lagrangian immersion in $\mathbb{CP}^2$. This is a soliton equation and its doubly periodic solutions are all of finite gap type: this fact can be deduced from [2] by recognising (1) as the affine $\mathfrak{a}_2^{(2)}$ Toda equations. It seems to have been first studied in this particular form by Sharipov [21] while he was studying “zero mean curvature, complex normal” tori in $S^5 \subset \mathbb{C}^3$. These are, one quickly realises, none other than the minimal Legendrian tori. Sharipov studies the finite gap solutions and gives fairly detailed formulae for the immersion in terms of Riemann θ -functions although he spares no room for examples. In particular, he points out that the spectral curve must possess an extra holomorphic involution. A few years later, Castro & Urbano [6] solved a reduction of (1) to an o.d.e while constructing S^1 -equivariant minimal Lagrangian tori in $\mathbb{CP}^2$. Haskins [11] performed a similar computation to find S^1 -equivariant minimal Legendrian tori in S^5 . Very recently, and following on from Sharipov, Ma & Ma [15] gave explicit formulae for the minimal Lagrangian maps into $\mathbb{CP}^2$ of finite type using Riemann θ -functions, although they do not seem to have noticed that their formulae arise from Sharipov's by Hopf projection $S^5 \rightarrow \mathbb{CP}^2$. Finally, Joyce [13] has given equivariant and non-equivariant minimal Lagrangian tori in $\mathbb{CP}^2$. His immersions are all given in terms of elliptic functions.

These results can be seen from a broader perspective, for there is a twistorial picture which will hold for cones over a surface of any topological type. In section three I describe two different but isomorphic primitive harmonic maps which determine a given special Lagrangian cone in $\mathbb{C}^3$. First, the Gauss map of the cone itself factors through a harmonic surface which admits a primitive lift into SU_3/SO_2 . Second, by applying the twistor theory of [2, 4] for the minimal surfaces in $\mathbb{CP}^2$, we construct a primitive lift into SU_3/S^1 . We view this as an outer 6-symmetric space, an S^1 -bundle over the 3-symmetric flag manifold which occurs in [2, 4]. The extra property of being Lagrangian imposes an extra involutive symmetry, corresponding to the outer involution with which we pass from $\mathfrak{a}_2$ to $\mathfrak{a}_2^{(2)}$.

In section four we restrict our attention to cones over tori. It turns out, due to a result of Bolton et al. [2], that every minimal Lagrangian torus in $\mathbb{CP}^2$ lifts to a minimal Legendrian torus in S^5 and thus there is a one-to-one correspondence between minimal Lagrangian tori in $\mathbb{CP}^2$ and special Lagrangian cones in $\mathbb{C}^3$ whose link is a torus. I recall how to construct minimal tori in $\mathbb{CP}^2$ and adapt this to the case of Lagrangian tori: I have tried to do this quickly without leaving out important details. In short, the spectral curve X of [16, 17] must be a double

cover $X \rightarrow Y$ of another curve, branched only at the unique zero P_0 and pole P_∞ of the degree three spectral parameter $\lambda : X \rightarrow \hat{\mathbb{C}}$. This means X has even genus $2p$, $p = \text{genus}(Y)$, and its Jacobian is isogenous to $Jac(Y) \times Prym(X, Y)$ (i.e. there is a homomorphism of $Jac(Y) \times Prym(X, Y)$ onto $Jac(X)$ with finite kernel). The line bundle $\mathcal{L}$ lies in a translate of $Prym(X, Y)$ and we can obtain families of minimal tori by moving $\mathcal{L}$ along this translate. A parameter count suggests that minimal Lagrangian tori exist for any spectral genus $2p$: in that case there will be large (real $(p - 2)$ -dimensional) continuous families of these obtained by moving along the Prym variety.

To finish we consider the case where the spectral curve has genus four; i.e., $p = 2$. Then $Prym(X, Y)$ is two dimensional and this forces the solutions of Tzitzéica's equation to be doubly periodic (although this is not sufficient to ensure the minimal map closes into a 2-torus). There are two types of spectral curves which can arise: I show that for one type $Prym(X, Y)$ is isogenous to a product of elliptic curves. This means the corresponding minimal maps may be constructed using elliptic functions. It is examples of this type which have been found by Joyce [13].

Acknowledgements. I would like to thank Dominic Joyce for sharing his results with me at an early stage and for encouraging me to think about them. I also thank Mark Haskins for his comments on the first draft.

Notation. For vectors $v_1, \dots, v_{n+1} \in \mathbb{C}^{n+1}$ we will use $v_1 \wedge \dots \wedge v_{n+1}$ to denote the *real* oriented $(n + 1)$ -plane with span $Sp_{\mathbb{R}}\{v_1, \dots, v_{n+1}\}$. Also, $(v_1, \dots, v_{n+1})$ will denote the matrix with j -th column v_j . For any Lie group G its connected component of the identity will be denoted G^0 . Finally, set $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$.

2. PRELIMINARIES.

A real oriented $(n + 1)$ -plane $V \subset \mathbb{C}^{n+1}$ is special Lagrangian if it lies in the SU_{n+1} orbit of the $(n + 1)$ -plane $e_1 \wedge \dots \wedge e_{n+1}$ where $\{e_j\}$ is the standard unitary basis for $\mathbb{C}^{n+1}$. Let SL denote the set of all these, then $SL \cong SU_{n+1}/SO_{n+1}$. Since $U_{n+1} \subset SO_{2n+2}$ is the subgroup of symplectic isometries of $\mathbb{C}^{n+1}$, equipped with its standard Kähler form and Riemannian metric, every element of SL is certainly Lagrangian. More generally we say V is special Lagrangian with phase $\theta \in [0, 2\pi)$ if

$$V \in SL_\theta = \{g(e_1 \wedge \dots \wedge e_{n+1}) : g \in U_{n+1}, \det(g) = e^{i\theta}\}.$$

A real oriented $(n + 1)$ -submanifold $N \subset \mathbb{C}^{n+1}$ is special Lagrangian if its Gauss map $\gamma_N : N \rightarrow Gr_{\mathbb{R}}(n + 1, \mathbb{C}^{n+1})$ takes values in SL . The importance of this property lies in the following two results of Harvey & Lawson [12]. First, if N is special Lagrangian then it is calibrated by the real $(n + 1)$ -form

$$\Omega_0 = \text{Re}(dz^1 \wedge \dots \wedge dz^{n+1})$$

and is therefore both minimal and Lagrangian. Conversely, if N is a connected minimal Lagrangian $(n + 1)$ -fold then it is congruent to a special Lagrangian $(n + 1)$ -fold. In particular, every minimal Lagrangian N is special Lagrangian with phase θ for some constant $\theta \in [0, 2\pi)$.

Let $\hat{\pi} : \mathbb{C}^{n+1} \setminus \{0\} \rightarrow S^{2n+1}$ denote the projection along rays to the unit sphere $S^{2n+1} \subset \mathbb{C}^{n+1}$. We will say $N \subset \mathbb{C}^{n+1}$ is a (regular) cone if $N = \hat{\pi}^{-1}(M) \cup \{0\}$ for some connected, oriented submanifold $M \subset S^{2n+1}$ and write $N = C(M)$. It follows that $M = C(M) \cap S^{2n+1}$ and M is called the link of the cone $C(M)$. Since $C(M)$ is usually singular at the origin we will say $C(M)$ is minimal whenever $C(M) \setminus \{0\}$ is minimal. It is obvious that $C(M)$ is Lagrangian whenever M is Legendrian for the contact structure S^{2n+1} inherits from $\mathbb{C}^{n+1}$. Moreover, since $\hat{\pi}$ is a Riemannian submersion with geodesic fibres, $C(M)$ is minimal if and only if M is minimal. We deduce (as did Haskins [11]) that there is a one-to-one correspondence between minimal Lagrangian $(n+1)$ -cones in $\mathbb{C}^{n+1}$ and minimal Legendrian n -folds in S^{2n+1} .

Now let $\pi : S^{2n+1} \rightarrow \mathbb{CP}^n$ be the Hopf fibration. Again, this is a Riemannian submersion with geodesic fibres, so if $M \subset S^{2n+1}$ is minimal then so is $\pi(M)$. Further, the pullback $\pi^*\omega$ of the Fubini-Study Kähler form is, up to a constant scalar, the curvature for the natural metric connection carried by the Hopf bundle. Moreover, the horizontal distribution is the standard contact structure on S^{2n+1} ; i.e., a Legendrian submanifold of S^{2n+1} is horizontal for π . Therefore every minimal Legendrian immersion $f : M \rightarrow S^{2n+1}$ induces a minimal Lagrangian immersion $\varphi = \pi \circ f : M \rightarrow \mathbb{CP}^n$. Conversely, a minimal Lagrangian $\varphi : M \rightarrow \mathbb{CP}^n$ has $\varphi^*\omega = 0$ and therefore has a horizontal lift of its universal cover $f : \tilde{M} \rightarrow S^{2n+1}$. It is a straightforward calculation to show that: (i) since f is horizontal the tension fields satisfy $\pi_*\tau(f) = \tau(\varphi)$ (see, for example, [8, p 20]), and; (ii) $\tau(f)$ is horizontal (using [20, §2]). Therefore f is minimal and, of course, Legendrian. So for simply connected manifolds there is, up to congruence, a one-to-one correspondence between minimal Lagrangian immersions in $\mathbb{CP}^n$ and minimal Legendrian immersions in S^{2n+1} .

3. TWISTOR THEORY FOR SPECIAL LAGRANGIAN CONES IN $\mathbb{C}^3$.

Let us now fix a special Lagrangian cone $C(M) \subset \mathbb{C}^3$ with link $f : M \rightarrow S^5$ and the corresponding minimal Lagrangian $\varphi : M \rightarrow \mathbb{CP}^2$. Our aim is to show that any one of these maps is characterised by the existence of a primitive harmonic lift of M into a certain 6-symmetric space. Let us first recall (from [4, §1]) the idea of a primitive harmonic map into a k -symmetric space.

A k -symmetric space is a reductive homogeneous space G/K for a compact semisimple Lie group G in which the isotropy group K is the subgroup of fixed points of an order k automorphism σ on G (or, more generally, we can take K to be a closed subgroup of the fixed point group which shares its Lie algebra). This automorphism induces an automorphism on $\mathfrak{g}$, the Lie algebra of G , which we will also call σ , with subalgebra of fixed points $\mathfrak{k}$, the Lie algebra of K . The complexification $\mathfrak{g}^{\mathbb{C}}$ decomposes into the direct sum of eigenspaces $\mathfrak{g}_j$ for eigenvalues ϵ^j where ϵ is a primitive k -th root of unity. The reductive decomposition of $\mathfrak{g}$ is characterised by

$$\mathfrak{g} = \mathfrak{k} + \mathfrak{p}; \quad \mathfrak{k}^{\mathbb{C}} = \mathfrak{g}_0, \quad \mathfrak{p}^{\mathbb{C}} = \sum_{j \neq 0} \mathfrak{g}_j.$$

Clearly K acts by adjoint action on each $\mathfrak{g}_j$. We recall that $T(G/K)$ is naturally isomorphic to $[\mathfrak{p}] = G \times_K \mathfrak{p}$ (the quotient of $G \times \mathfrak{p}$ by right K -action). We can identify this with a subbundle of $G/K \times \mathfrak{g}$ so that fibrewise the isomorphism is

$$[\mathfrak{p}]_q \rightarrow T_q(G/K); \quad X \mapsto \frac{d}{dt}(e^{tX} \cdot q)|_{t=0}.$$

In particular $[\mathfrak{p}]_q = \text{Ad } g \cdot \mathfrak{p}$ whenever $q = gK$. The complexification $T^{\mathbb{C}}(G/K)$ possesses subbundles $[\mathfrak{g}_j]$ for $j \neq 0$. For $k \geq 3$, a smoothly immersed surface $\psi : M \rightarrow G/K$ is said to be primitive whenever

$$(2) \quad d\psi^{1,0} : TM^{1,0} \rightarrow [\mathfrak{g}_1].$$

This is a weakened version of the J -holomorphic condition which one finds in Riemannian twistor theory. Let $F \in \text{End}(T^{\mathbb{C}}G/K)$ be the bundle endomorphism satisfying $F^3 = -F$ and having $[\mathfrak{g}_1]$ as its bundle of i -eigenspaces, $[\mathfrak{g}_{k-1}]$ as its bundle of $-i$ -eigenspaces and kernel $\oplus_{j=2}^{k-2} [\mathfrak{g}_j]$. Then F is one of the horizontal f -structures studied by Black [1, §6]. The condition (2) means ψ is f -holomorphic with respect to this f -structure. By results of Black [1, Thm 4.3, Lemma 5.3] we have: (1) every primitive map is harmonic (for any G -invariant metric on G/K for which $[\mathfrak{g}_1]$ is isotropic); (2) if $H \subset G$ is any closed subgroup containing K then post-composition of any primitive map ψ with the homogeneous projection $G/K \rightarrow G/H$ again yields a harmonic map.

Given the special Lagrangian cone $C(M)$ we will find two primitive harmonic maps. The first, $\psi_1 : M \rightarrow SU_3/SO_2$, arises from the Gauss map of $C(M)$. The second, $\psi_2 : M \rightarrow SU_3/S^1$, arises by considering how the Lagrangian condition interacts with the primitive lift assigned by [2, 4] to any minimal surface in $\mathbb{CP}^2$. We will show that these are basically the same map.

3.1. The Gauss map of $C(M)$. Let $\gamma_{C(M)} : C(M) \setminus \{0\} \rightarrow SL$ be the Gauss map of $C(M)$. Note that since $C(M)$ is minimal $\gamma_{C(M)}$ is harmonic by the Ruh-Vilms theorem (see, for example, [8, p 18]). Since $C(M)$ is a cone $\gamma_{C(M)}$ factors through M . Indeed, it is easy to see that if $\gamma_M : M \rightarrow Gr_{\mathbb{R}}(2, \mathbb{C}^3)$ denotes the Gauss map of the link $f : M \rightarrow S^5$ then $\gamma_{C(M)}$ is projection onto M followed by

$$(3) \quad f \wedge \gamma_M : M \rightarrow SL \cong SU_3/SO_3.$$

This too is harmonic since the projection is Riemannian with geodesic fibres.

Now set

$$FL_1 = \{(v, V) \in S^5 \times SL : v \in V\} = \{(ge_1, g(e_1 \wedge e_2 \wedge e_3)) : g \in SU_3\}.$$

Clearly $FL_1 \cong SU_3/K_1$ where $K_1 = \{g \in SO_3 : ge_1 = e_1\} \cong SO_2$ and the two natural projections

$$\pi_1 : SU_3/K_1 \rightarrow SU_3/SU_2 \cong S^5; \quad \pi_2 : SU_3/K_1 \rightarrow SU_3/SO_3 \cong SL$$

are homogeneous projections. The map (3) obviously possesses a lift

$$\psi_1 : M \rightarrow FL_1; \quad \psi_1 = (f, f \wedge \gamma_M).$$

We can view FL_1 as a 6-symmetric space as follows. Let $\hat{\sigma} \in \text{Aut}(SU_3)$ be given by

$$\hat{\sigma}(g) = Rg^{-1t}R^{-1},$$

where $R \in SO_3$ is the rotation about the e_1 axis through angle $\pi/3$. Then $\hat{\sigma}$ is an outer automorphism of order six with subgroup of fixed points K_1 .

Proposition 1. *The map $\psi_1 : M \rightarrow FL_1$ is primitive harmonic. Conversely, any such map projects by π_1 to the Gauss map of an immersed special Lagrangian cone $\kappa : C(M) \rightarrow \mathbb{C}^3$ and by π_2 to a minimal Legendrian map $f : M \rightarrow S^5$.*

The proof of this will follow once we have proved a similar result for the primitive lift of the minimal Lagrangian map $\varphi : M \rightarrow \mathbb{CP}^2$.

3.2. Minimal Lagrangian surfaces in $\mathbb{CP}^2$. As we noted above, every minimal Lagrangian $\varphi : M \rightarrow \mathbb{CP}^2$ lifts (locally or on the universal cover) horizontally to a minimal Legendrian map into S^5 . For simplicity let us assume this lift is global $f : M \rightarrow S^5$; i.e., φ arises from a special Lagrangian cone in $\mathbb{C}^3$. We will use ideas from [3, 4] to lift φ into the manifold

$$FL_2 = \{(w, W) \in S^5 \times Gr_{\mathbb{C}}(2, \mathbb{C}^3) : w \in W\}.$$

The motivation for this is that [2, 4] have already shown that any minimal surface in $\mathbb{CP}^2$ lifts to a primitive map into the flag manifold

$$Fl_2 = \{(W_1, W_2) \in \mathbb{CP}^2 \times Gr_{\mathbb{C}}(2, \mathbb{C}^3) : W_1 \subset W_2\}.$$

We will show that the extra symmetry of being Lagrangian allows us to lift to the S^1 -bundle $pr_2 : FL_2 \rightarrow Fl_2$ and the lift will also be primitive. If we think of Fl_2 as the 3-symmetric space for the Coxeter-Killing automorphism ν (whose definition will be recalled shortly) then FL_2 is the 6-symmetric space for an automorphism σ which is the product of ν and a commuting involution μ .

To begin, let $L \rightarrow \mathbb{CP}^2$ denote the tautological line bundle, then φ determines (and is determined by) $\ell_0 = \varphi^*L \subset M \times \mathbb{C}^3$. Any smooth subbundle $\ell \subset M \times \mathbb{C}^3$ can be equipped with a holomorphic structure for which a local section is holomorphic whenever $\pi_{\ell}(\partial s / \partial \bar{z}) = 0$, where π_{ℓ} is the orthogonal projection onto ℓ and z is a local complex coordinate on M . Recall that φ is harmonic precisely when the map

$$d\varphi\left(\frac{\partial}{\partial z}\right) : \ell_0 \rightarrow \ell_0^{\perp}; \quad s \mapsto \pi_{\ell_0}^{\perp}\left(\frac{\partial s}{\partial z}\right),$$

is holomorphic. It is well known that in this case the new line bundles $\ell_1, \ell_2, \dots$ defined by

$$(4) \quad \ell_{j+1} = \pi_{\ell_j}^{\perp}\left(\frac{\partial}{\partial z}\ell_j\right),$$

each determine a harmonic map $\varphi_j : M \rightarrow \mathbb{CP}^n$ for which $\ell_j = \varphi_j^*L$. The sequence $\ell_0, \ell_1, \dots$ is called the harmonic sequence (see eg. [3]). The extra condition of conformality is equivalent to $\ell_2 \perp \ell_0$. Thus from a minimal surface

in $\mathbb{CP}^2$ we obtain an orthogonal harmonic sequence ℓ_0, ℓ_1, ℓ_2 which decomposes $M \times \mathbb{C}^3$. The primitive harmonic lift of φ is the map

$$\psi : M \rightarrow FL_2; \psi = (\ell_0, \ell_0 \oplus \ell_1).$$

Now, since φ is minimal Lagrangian with the lift $f : M \rightarrow S^5$, we can define

$$\psi_2 : M \rightarrow FL_2; \psi_2 = (f, \ell_0 \oplus \ell_1).$$

We will show that this is primitive harmonic with respect to a particular 6-symmetric space structure on FL_2 .

First, notice that FL_2 is the SU_3 -orbit of (w_0, W_0) , where

$$w_0 = e_1, W_0 = Sp_{\mathbb{C}}\{e_1, e_2\}.$$

The isotropy group is $K_2 = \{\text{diag}(1, a, a^{-1}) \in SU_3\}$. Therefore $FL_2 \cong SU_3/K_2$ and it fibres over the flag manifold $Fl_2 \cong SU_3/T$ (where T is the maximal torus of diagonal matrices) with homogeneous projection

$$pr_2 : FL_2 \rightarrow Fl_2; (w, W) \mapsto (Sp_{\mathbb{C}}\{w\}, W).$$

We view FL_2 as the 6-symmetric space corresponding to the order six automorphism $\sigma \in \text{Aut}(SU_3)$ given by the product $\sigma = \mu\nu$ where: (i) ν is the Coxeter-Killing automorphism $\nu(g) = SgS^{-1}$ where $S = \text{diag}(1, \epsilon, \epsilon^2)$ for $\epsilon = \exp(2\pi i/3)$; (ii) μ is the involution $\mu(g) = Tg^{-1}T^{-1}$ where

$$(5) \quad Te_1 = e_1, Te_2 = e_3, Te_3 = e_2.$$

These two automorphisms commute and so σ is well-defined, with subgroup of fixed points K_2 . On $\mathfrak{g} = \mathfrak{su}_3$ we take the eigenvalues of σ to be $\{(-\epsilon)^j : j = 0, \dots, 5\}$: this makes it easier to work with the eigenspaces $\mathfrak{g}_j$ of σ , by observing that

$$(6) \quad \mathfrak{g}_j = \mathfrak{g}_{j \bmod 2}^{\mu} \cap \mathfrak{g}_{j \bmod 3}^{\nu}.$$

In particular

$$\mathfrak{g}_1 = \{X \in \mathfrak{g}^{\mathbb{C}} : X = \begin{pmatrix} 0 & 0 & \alpha \\ \alpha & 0 & 0 \\ 0 & \beta & 0 \end{pmatrix}, \alpha, \beta \in \mathbb{C}\}$$

Let $\mathfrak{g} = \mathfrak{k}_2 \oplus \mathfrak{p}_2$ be the corresponding reductive splitting and make the usual identification

$$(7) \quad T^{\mathbb{C}}FL_2 \cong [\mathfrak{p}_2^{\mathbb{C}}] = SU_3 \times_{K_2} \mathfrak{p}_2^{\mathbb{C}}.$$

The primitive distribution is the subbundle $[\mathfrak{g}_1]$ whose fibres can be described as

$$[\mathfrak{g}_1]_q = \text{Ad}g \cdot \mathfrak{g}_1 \subset \mathfrak{g}^{\mathbb{C}}; \text{ at } q = gK_2.$$

Proposition 2. *Let $\varphi : M \rightarrow \mathbb{CP}^2$ be minimal Lagrangian with horizontal lift $f : M \rightarrow S^5$. Then we can choose f so that the lift $\psi_2 : M \rightarrow FL_2$ is primitive harmonic; i.e., $d\psi_2^{1,0} \in \Omega_M^{1,0} \otimes \psi_2^*[\mathfrak{g}_1]$ for the 6-symmetric space structure described above. Conversely, any primitive harmonic ψ_2 projects by $pr_1 : FL_2 \rightarrow S^5$ to a minimal Legendrian surface in S^5 and thence by the Hopf map to a minimal Lagrangian surface in $\mathbb{CP}^2$.*

Remark. This special choice of horizontal lift f is an artefact of our definition of FL_2 . In the next subsection I will show that any choice of f gives rise to some primitive lift into an isomorphic 6-symmetric space.

Proof. We can move f by an isometry of the form $f \mapsto e^{i\theta}f$ so as to ensure that the Lagrangian map $f \wedge \gamma_M$ is special with phase $3\pi/2$. In terms of a local complex conformal coordinate $z = x + iy$ we have

$$\det(f, \frac{1}{|f_x|}f_x, \frac{1}{|f_y|}f_y) = -i.$$

Now we can choose a local frame F for ψ_2 by taking

$$(8) \quad F = (f, \frac{1}{|f_z|}f_z, \frac{-1}{|f_{\bar{z}}|}f_{\bar{z}}) = (f_0, f_1, f_2).$$

Now, f is a horizontal (and therefore holomorphic) section of ℓ_0 , and since φ is conformal we have

$$\langle f_z, f \rangle = 0 = \langle f_{\bar{z}}, f \rangle, \quad \langle f_z, f_{\bar{z}} \rangle = 0.$$

Therefore each f_j is a local section of ℓ_j and F is unitary. Further,

$$(9) \quad (f, \frac{1}{|f_x|}f_x, \frac{1}{|f_y|}f_y) = FU \text{ for } U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1/\sqrt{2} & -i/\sqrt{2} \\ 0 & -1/\sqrt{2} & -i/\sqrt{2} \end{pmatrix}.$$

Since $\det(U) = -i$ it follows that F is special unitary. Now recall that using the identification (7) we identify

$$d\psi_2(\partial/\partial z) = \text{Ad}F \cdot (F^{-1} \frac{\partial F}{\partial z})_{\mathfrak{p}_2^{\mathbb{C}}}.$$

A straightforward calculation gives

$$(10) \quad (F^{-1} \frac{\partial F}{\partial z})_{\mathfrak{p}_2^{\mathbb{C}}} = \begin{pmatrix} 0 & 0 & |f_{\bar{z}}| \\ |f_z| & 0 & 0 \\ 0 & Q|f_z|^{-2} & 0 \end{pmatrix}$$

where $Q = \langle f_{zzz}, f \rangle$. But now we observe that $\langle f_{\bar{z}z}, f \rangle = \langle f_{z\bar{z}}, f \rangle$ implies that $|f_z| = |f_{\bar{z}}|$ and therefore $d\psi_2^{1,0}$ takes values in the primitive distribution. The converse statement in the proposition follows at once from the result due to Black [1] mentioned earlier. $\square$

A standard argument (see, for example, [2, p 125]) shows that the cubic form $\det(d\psi_2^{1,0}) = Qdz^3$ is holomorphic on M , since ψ_2 is harmonic for the SU_3 -invariant metric. Indeed, Qdz^3 is globally defined even if the lift $f : M \rightarrow S^5$ is not global since it is easy to see that $d\psi_2$ is the pullback of $d\psi$ under $FL_2 \rightarrow Fl_2$, and $\psi : M \rightarrow Fl_2$ exists as a global lift for any minimal surface in $\mathbb{CP}^2$. The form Qdz^3 is the Hopf differential of φ : its zeroes represent the higher order singularities of ψ .

3.3. The isomorphism between FL_1 and FL_2 . In the previous section we found a lift ψ_2 of the minimal Lagrangian map φ which corresponded to a special Lagrangian cone with phase $3\pi/2$. This phase factor can be avoided if we perform the following sleight of hand. We identify FL_2 with N/K_2 where

$$N = \{g \in U_3 : \det(g) = i\} = \{U^{-1}g : g \in SU_3\},$$

and U is the matrix defined in (9). The implicit identification $N/K_2 \cong SU_3/K_2$ allows us to identify $T^{\mathbb{C}}N/K_2$ with $[\mathfrak{p}_2^{\mathbb{C}}]$ so that the primitive distribution has fibre

$$[\mathfrak{g}_1]_q = \text{Ad}g \cdot \mathfrak{g}_1 \subset \mathfrak{g}^{\mathbb{C}}, \text{ at } q = U^{-1}gK_2.$$

Now if we take $f : M \rightarrow S^5$ to be a special Legendrian lift of φ , with phase zero, the local frame F in (8) has $\det(F) = i$ so that $\psi_2 = (f, \ell_0 \oplus \ell_1)$ is a primitive harmonic lift.

With this slightly altered point of view there is a natural isomorphism $\Phi : FL_2 \cong FL_1$ for which $\Phi_*[\mathfrak{g}_1]$ is the primitive distribution for FL_1 and for which $\psi_1 = \Phi \circ \psi_2$. Define

$$\Phi : FL_2 \rightarrow FL_1; (w, W) \mapsto (v, V),$$

by $v = w$ and $V = v_0 \wedge v_1 \wedge v_2$ where

$$(v, v_1, v_2) = (w, \frac{1}{\sqrt{2}}(w_1 - w_2), \frac{-i}{\sqrt{2}}(w_1 + w_2)) = (w, w_1, w_2)U.$$

Here w, w_1, w_2 is any unitary basis of $\mathbb{C}^3$ adapted to the flag $Sp_{\mathbb{C}}\{w\} \subset W \subset \mathbb{C}^3$ and with $\det(w, w_1, w_2) = i$. It is easy to check that Φ is independent of which basis we choose and

$$\det(v, v_1, v_2) = 1.$$

Further, since U is unitary the real 3-plane V is Lagrangian, whence $V \in SL$.

Proposition 3. *The isomorphism $\Phi : FL_2 \rightarrow FL_1$ has the properties: (i) $\Phi_*[\mathfrak{g}_1] = [\hat{\mathfrak{g}}_1]$ where $[\hat{\mathfrak{g}}_1]$ is the primitive distribution for the 6-symmetric space structure on FL_1 determined by $\hat{\sigma}$ given earlier; (ii) $\psi_1 = \Phi \circ \psi_2$. Therefore ψ_1 is primitive harmonic whenever ψ_2 is.*

Proof. Clearly $\psi_1 = \Phi \circ \psi_2$ by (9). Now let us consider $\Phi_* : T^{\mathbb{C}}FL_2 \rightarrow T^{\mathbb{C}}FL_1$. The identification of FL_1 with SU_3/K_1 gives the usual isomorphism

$$T_p^{\mathbb{C}}FL_1 \cong \text{Ad}g \cdot \mathfrak{p}_1^{\mathbb{C}} \subset \mathfrak{g}^{\mathbb{C}}, \text{ at } p = gK_1.$$

We can represent Φ by

$$\Phi : N/K_2 \rightarrow SU_3/K_1; U^{-1}gK_2 \rightarrow U^{-1}gUK_2, \det(g) = 1,$$

(the reader can check that for any $k \in K_2$ we have $U^{-1}kU \in K_1$). Therefore

$$\Phi_* : [\mathfrak{p}_2^{\mathbb{C}}] \rightarrow [\mathfrak{p}_1^{\mathbb{C}}]; \Phi_*(X) = \text{Ad}U^{-1} \cdot X.$$

Now consider $X \in [\mathfrak{g}_1]_q$. We can write $X = \text{Ad}g \cdot Y$ where $Y \in \mathfrak{g}_1$; i.e.,

$$-\text{Ad}TS \cdot Y^t = -\epsilon Y.$$

Then $\Phi_*(X) = \text{Ad}U^{-1}g \cdot Y$ so

$$\Phi_*(X) = \text{Ad}U^{-1}gU \cdot \hat{Y}, \text{ for } \hat{Y} = \text{Ad}U^{-1} \cdot Y.$$

Therefore

$$\text{Ad}TS \cdot (\text{Ad}U \cdot \hat{Y})^t = \epsilon \text{Ad}U \cdot \hat{Y}$$

hence $-\text{Ad}R \cdot \hat{Y}^t = -\epsilon \hat{Y}$, where

$$R = U^{-1}TSU^{-1t} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\pi/3) & \sin(\pi/3) \\ 0 & -\sin(\pi/3) & \cos(\pi/3) \end{pmatrix}.$$

But FL_1 has the 6-symmetric space structure corresponding to the automorphism $\hat{\sigma}(g) = Rg^{-1t}R$ of SU_3 , for which the primitive distribution is determined by

$$\hat{\mathfrak{g}}_1 = \{X \in \mathfrak{g}^{\mathbb{C}} : -\text{Ad}R \cdot X^t = -\epsilon X\}.$$

Thus $\Phi_*[\mathfrak{g}_1] = [\hat{\mathfrak{g}}_1]$. □

Proposition 1 follows now from proposition 2.

4. MINIMAL LAGRANGIAN TORI IN $\mathbb{CP}^2$.

Let us now suppose we have a minimal Lagrangian torus $\varphi : T^2 \rightarrow \mathbb{CP}^2$. In this case the Hopf differential Q must be constant. Either $Q = 0$, in which case φ is superminimal, or $Q \neq 0$ and φ is superconformal. One knows from [3, Thm 3.6] that a superminimal Lagrangian surface must have image congruent to $\mathbb{RP}^2$: the corresponding special Lagrangian cone is $\mathbb{R}^3 \subset \mathbb{C}^3$. So we may assume φ is superconformal and therefore possesses a *Toda frame*. This means (see [2, §2]) there is a frame $F : \mathbb{R}^2 \rightarrow SU_3$ satisfying, in an appropriate coordinate z , the equation

$$(11) \quad F^{-1} \frac{\partial F}{\partial z} = \begin{pmatrix} s_0^{-1} \partial s_0 / \partial z & 0 & s_0 s_2^{-1} \\ s_1 s_0^{-1} & s_1^{-1} \partial s_1 / \partial z & 0 \\ 0 & s_2 s_1^{-1} & s_2^{-1} \partial s_2 / \partial z \end{pmatrix},$$

where $s_j : T^2 \rightarrow \mathbb{R}^+$ satisfy the affine $\mathfrak{a}_2$ Toda equations

$$(12) \quad \frac{\partial^2}{\partial z \partial \bar{z}} \log(s_j^2) = s_{j+1}^2 s_j^{-2} - s_j^2 s_{j-1}^{-2}, \quad j \in \mathbb{Z}_3.$$

Recall that the latter are simply the integrability conditions for the existence of F . Moreover, it was shown in [2, Thm 2.5] that Toda frames are essentially uniquely determined by this property and they are also double periodic, to wit, $F : \tilde{T}^2 \rightarrow SU_3$ where $\tilde{T}^2 \rightarrow T^2$ is a covering torus with covering group either 1 or $\mathbb{Z}_3$. Using these facts let us now prove:

Proposition 4. *Every minimal Lagrangian torus $\varphi : T^2 \rightarrow \mathbb{CP}^2$ has a doubly periodic primitive lift $\psi_2 : \tilde{T}^2 \rightarrow FL_2$ where $\tilde{T}^2 \rightarrow T^2$ has covering group either 1 or $\mathbb{Z}_3$. Consequently φ determines a minimal Legendrian $f : \tilde{T}^2 \rightarrow S^5$ and an immersed special Lagrangian cone $\kappa : C(\tilde{T}^2) \rightarrow \mathbb{C}^3$.*

Proof. It suffices to show that, with an appropriate choice of complex coordinate z on $\mathbb{R}^2$ the frame $F = (f_0, f_1, f_2)$ given by (8) satisfies the Toda frame equations. Since the Hopf differential Qdz^3 is globally holomorphic on T^2 we can choose z such that $Q = \langle f_{zzz}, f \rangle = 1$. Now define $s_j : \mathbb{R}^2 \rightarrow \mathbb{R}^+$ by

$s_0 = 1, s_1 = |f_z|, s_2 = 1/|f_z|$. The equation (10) shows that F satisfies (11) provided we can show that

$$\langle \frac{\partial f_j}{\partial z}, f_j \rangle = s_j^{-1} \frac{\partial}{\partial z} s_j, \quad j \in \mathbb{Z}_3.$$

The reader is left to check that these equations are true using the identities $f_1 = s_1^{-1} f_z, f_2 = -s_2 f_{\bar{z}}$ and

$$\langle f_z, f \rangle = 0, \quad \langle f_{z\bar{z}}, f_z \rangle = 0 = \langle f_{z\bar{z}}, f_{\bar{z}} \rangle.$$

The last pair of equations follows from the fact that $f_z = d\varphi(f \otimes \partial/\partial z)$ is a holomorphic section of ℓ_1 while $f_{\bar{z}} = d\varphi(f \otimes \partial/\partial \bar{z})$ is an anti-holomorphic section of ℓ_2 , since φ is harmonic. Here we are interpreting $d\varphi$ as a 1-form with values in $\varphi^{-1}T\mathbb{CP}^2 \cong \text{Hom}(\ell_0, \ell_0^\perp)$ (cf. [3]). $\square$

4.1. The spectral data for a minimal Lagrangian torus. Recall from [2, 4] that for every minimal torus $\varphi : T^2 \rightarrow \mathbb{CP}^2$ its primitive lift $\psi : T^2 \rightarrow Fl_2$ is of semisimple finite type. This ultimately means that φ is determined, uniquely up to congruence, by its spectral data $(X, \lambda, \mathcal{L})$ [16]. Here I will describe what this data is for a minimal Lagrangian torus: basically, the extra condition of being Lagrangian equips X with a holomorphic involution which λ and $\mathcal{L}$ must also respect.

As above, let $F = (f_0, f_1, f_2)$ be the Toda frame for a minimal Lagrangian torus $\varphi : M \rightarrow \mathbb{CP}^2$. Set $\alpha = F^{-1}dF$, and write $\alpha = \alpha_{\mathfrak{k}} + \alpha_{\mathfrak{p}}$ for the components in the reductive decomposition of $\mathfrak{g}$. Since $s_0 = 1, s_2 = s_1^{-1}$ these components satisfy

$$\alpha_{\mathfrak{k}}\left(\frac{\partial}{\partial z}\right) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & s_1^{-1}\partial s_1/\partial z & 0 \\ 0 & 0 & -s_1^{-1}\partial s_1/\partial z \end{pmatrix}, \quad \alpha_{\mathfrak{p}}\left(\frac{\partial}{\partial z}\right) = \begin{pmatrix} 0 & 0 & s_1 \\ s_1 & 0 & 0 \\ 0 & s_1^{-2} & 0 \end{pmatrix}.$$

Further, $s_1 : T^2 \rightarrow \mathbb{R}^+$ satisfies the affine $\mathfrak{a}_2^{(2)}$ Toda equations

$$(13) \quad \frac{\partial^2}{\partial z \partial \bar{z}} \log(s_1^2) = s_1^{-4} - s_1^2.$$

Notice that $s_1 = \|\partial\varphi/\partial z\|$ and therefore s_1^2 represents the conformal factor in the metric induced on T^2 by the immersion φ (cf. [6, §2]).

Now we recall from [5, §3] that the lift ψ_2 is harmonic if and only if any frame can be extended to an S^1 -family (or loop) of frames

$$F_\zeta : \mathbb{R}^2 \rightarrow \Lambda^\sigma SU_3 = \{g_\zeta : S^1 \rightarrow SU_3 | \sigma(g_\zeta) = g_{-\epsilon\zeta}\}.$$

This loop of frames is determined, up to initial condition, by

$$F_\zeta^{-1}dF_\zeta = \alpha_\zeta = \zeta\alpha_{\mathfrak{p}}^{1,0} + \alpha_{\mathfrak{k}} + \zeta^{-1}\alpha_{\mathfrak{p}}^{0,1}.$$

We note that α_ζ actually possesses three symmetries

$$\nu(\alpha_\zeta) = \alpha_{\epsilon\zeta}, \quad \mu(\alpha_\zeta) = \alpha_{-\zeta}, \quad -\alpha_\zeta^\dagger = \alpha_{\bar{\zeta}^{-1}},$$

(the first two can be deduced from (6)), which obliges F_ζ to have the same symmetries (for every initial condition with these symmetries).

We recall from [18, p 240] that there is a commutative algebra $\mathcal{A}$ of complexified polynomial Killing fields

$$\xi_\zeta : \mathbb{R}^2 \rightarrow \Lambda^\nu \mathfrak{gl}_3 = \{X_\zeta : S^1 \rightarrow \mathfrak{gl}_3(\mathbb{C}) | \nu(X_\zeta) = X_{\epsilon_\zeta}\}, \quad d\xi_\zeta = [\xi_\zeta, \alpha_\zeta],$$

where ξ_ζ is a Laurent polynomial in ζ (of bounded degree over $\mathbb{R}^2$). By the symmetries of α_ζ , this algebra admits two automorphisms

$$\mu_*(\xi_\zeta) = \mu(\xi_{-\zeta}), \quad \rho_*(\xi_\zeta) = -\bar{\xi}_{\zeta^{-1}}^t.$$

The algebra $\mathcal{A}$ determines the spectral curve X of φ , as the smooth completion of the the affine curve $\text{Spec}(\mathcal{A})$, together with a degree three cover $\lambda : X \rightarrow \hat{\mathbb{C}}$ with a single zero P_0 and pole P_∞ . This cover is the dual of $\mathbb{C}[\zeta^3 I] \subset \mathcal{A}$, where I is the identity matrix. The real involution ρ_* equips X with a real involution ρ covering $\lambda \mapsto \bar{\lambda}^{-1}$, for which ρ fixes every point over $|\lambda| = 1$, while μ_* induces a holomorphic involution μ on X which fixes only the points P_0, P_∞ (since it must cover $\lambda \mapsto -\lambda$). Inside $\mathcal{A}$ we have the subalgebra $\mathcal{A}^\mu = \{\xi_\zeta : \mu_*(\xi_\zeta) = \xi_\zeta\}$, which we might naturally think of as the polynomial Killing fields for ψ_2 .

Lemma 1. $\mathcal{A} = \mathcal{A}^\mu[\zeta^3 I] \cong \mathcal{A}^\mu[t]/(t^2 - \zeta^6 I)$, where $(t^2 - \zeta^6 I)$ denotes the ideal generated by this quadratic. Therefore the inclusion $\mathcal{A}^\mu \subset \mathcal{A}$ is dual to a double cover $\delta : X \rightarrow Y$, ramified at P_0 and P_∞ only, where Y is the smooth completion of $\text{Spec}(\mathcal{A}^\mu)$. The involution μ is the swapping of sheets for this cover. It follows that X has even genus $g = 2p$ where p is the genus of Y .

Proof. We can write each ξ_ζ as $A_\zeta + \zeta^3 B_\zeta$ where

$$A_\zeta = \xi_\zeta + \mu_*(\xi_\zeta), \quad B_\zeta = \zeta^{-3}(\xi_\zeta - \mu_*(\xi_\zeta)).$$

Then $A_\zeta, B_\zeta \in \mathcal{A}^\mu$ and we notice that $\mathbb{C}[\zeta^6 I] \subset \mathcal{A}^\mu$. It follows that $\mathcal{A} = \mathcal{A}^\mu[\zeta^3 I]$. Therefore X is the Riemann surface for the quadratic extension $t^2 - \zeta^6 I = 0$ over the smooth completion Y of $\text{Spec}(\mathcal{A}^\mu)$. This is clearly only ramified at P_0, P_∞ . $\square$

Next we recall from [18] that any minimal torus φ is determined, up to congruence, by the data $(X, \lambda, \mathcal{L})$ where $\mathcal{L}$ is a holomorphic line bundle over X of degree $g + 2$. To be precise, we identify T^2 with $\mathbb{C}/\Lambda$ for some lattice Λ and assign to $\varphi(z)$ the data $(X, \lambda, \mathcal{L}_z)$: this whole family is determined by the initial data $(X, \lambda, \mathcal{L}_0)$. Let us recall how this works. We will think of $\mathcal{L}_z$ as the dual to the eigenline bundle $\mathcal{E}_z$ over X , defined as follows. First, to each $\xi_\zeta \in \mathcal{A}$ we assign an untwisted polynomial Killing field

$$\hat{\xi}_{\zeta^3} = \text{Ad} \kappa_\zeta \cdot \xi_\zeta, \quad \kappa_\zeta = \text{diag}(1, \zeta^{-1}, \zeta^{-2}).$$

Since $\text{Ad} \kappa_{\epsilon_\zeta} = \text{Ad} \kappa_\zeta \circ \nu$ we deduce that this untwisted loop really is a function of $\lambda = \zeta^3$. This provides an isomorphic algebra $\hat{\mathcal{A}} = \{\hat{\xi}_\lambda : \xi_\zeta \in \mathcal{A}\}$. Notice that this untwisting has not changed the symmetry $\mu_* : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{A}}; \mu_*(\hat{\xi}_\lambda) = \mu(\hat{\xi}_{-\lambda})$. For each $z \in \mathbb{C}$ we can model X on the smooth completion X_z of the curve

$$X_z \setminus \{P_0, P_\infty\} \cong \{(\lambda, [v]) \in \mathbb{C} \setminus \{0\} \times \mathbb{P}^2 : \hat{\xi}_\lambda(z)[v] = [v], \forall \hat{\xi}_\lambda \in \hat{\mathcal{A}}\}.$$

It is also not hard to show that all these curves are isomorphic (cf. [9]) and the completion is obtained by adding the points $P_0 = (0, [0, 0, 1])$ and $P_\infty = (\infty, [1, 0, 0])$. For each z this embeds X in $\mathbb{P}^1 \times \mathbb{P}^2$ and $\mathcal{E}_z$ is the pullback of the

tautological bundle over $\mathbb{P}^2$; i.e., the bundle of common eigenlines for all $\hat{\xi}_\lambda(z)$. A necessary condition for $(X, \lambda, \mathcal{L}_z)$ to be the spectral data for a minimal torus is that $\mathcal{L}_z$ satisfies the reality condition

$$(14) \quad \overline{\rho_* \mathcal{L}_z} \otimes \mathcal{L}_z \cong \mathcal{O}_X(R)$$

where R is the divisor of ramification divisor of λ . [In fact slightly more must be true: this linear equivalence must be achieved by a function on X which is positive over the unit circle. This is always the case if, for example, $\mathcal{L}_z \cong \mathcal{O}_X(R_+) \otimes L$ where R_+ is the divisor of those ramification points lying over $|\lambda| < 1$ and L lies in the identity component of $\{L \in \text{Jac}(X) : \overline{\rho_* L} \cong L^{-1}\}$: see [17, §2.2].]

Lemma 2. *For a minimal Lagrangian torus in $\mathbb{CP}^2$ the data $(X, \lambda, \mathcal{L}_z)$ has the extra property that*

$$(15) \quad \mu_* \mathcal{L}_z \cong \mathcal{L}_z^{-1} \otimes \mathcal{O}_X(R).$$

The family of line bundles $L_z = \mathcal{L}_z \otimes \mathcal{L}_0^{-1}$ provides a real homomorphism $L : \mathbb{R}^2 \rightarrow P_{\mathbb{R}}(X, \mu)$, where

$$P_{\mathbb{R}}(X, \mu) = \{L \in \text{Prym}(X, \mu) : \overline{\rho_* L} \cong L^{-1}\}^0.$$

This homomorphism is uniquely determined by the property that

$$(16) \quad dL_0\left(\frac{\partial}{\partial z}\right) = d\mathcal{A}_{P_\infty}\left(\frac{\partial}{\partial \zeta^{-1}}\right)$$

where $\mathcal{A}_{P_\infty} : X \rightarrow \text{Jac}(X)$ is the Abel map with base point P_∞ .

Proof. We will prove the equivalent statement $\mu_* \mathcal{E}_z \cong \text{Hom}(\mathcal{E}_z, \mathcal{O}_X(-R))$. Let $U \subset X$ be any proper open subset with $\mu(U) = U$ and $\lambda^{-1}(\lambda(U)) = U$, so that $U \rightarrow \lambda(U)$ is a 3-sheeted cover. Since $\mathcal{E} \subset X \times \mathbb{C}^3$ any trivializing section v for $\mathcal{E}_z$ over U determines a matrix function V_λ by

$$V_{\lambda_0} = (v_{P_1}, v_{P_2}, v_{P_3}), \quad \lambda(P_j) = \lambda_0.$$

Note that $\det(V_\lambda) = 0$ iff λ is a branch point. Clearly $V_\lambda^{-1} \hat{\xi}_\lambda V_\lambda$ is diagonal for every $\hat{\xi}_\lambda \in \hat{\mathcal{A}}$: we will write

$$V_\lambda^{-1} \hat{\xi}_\lambda V_\lambda = D(\hat{\xi}_\lambda).$$

Now define $W_\lambda = \det(V_\lambda) T V_\lambda^{-1t}$, where T is the matrix from (5). Notice that W_λ is holomorphic on $\lambda(U)$: it is a constant matrix times the classical adjoint matrix for V_λ . Now we observe that

$$\begin{aligned} W_\lambda^{-1} \mu_*(\hat{\xi}_{-\lambda}) W_\lambda &= -V_\lambda^t T^2 \hat{\xi}_\lambda T^2 V_\lambda^{-1t} \\ &= -(V_\lambda^{-1} \hat{\xi}_\lambda V_\lambda)^t \\ &= -D(\hat{\xi}_\lambda). \end{aligned}$$

Since $\mu : X \rightarrow X; (\lambda, [v]) \mapsto (-\lambda, \mu[v])$ if we define $\mu_*(\hat{\xi}_{-\lambda})\mu[v] = \mu[v]$ we deduce that W_λ represents a section w of $\mu_* \mathcal{E}_z$ over U . Further, none of the columns of W_λ vanish on U so w is a trivialising section. On the other hand $W_\lambda^t T V_\lambda = \det(V_\lambda)$ implies that w is a section of $\text{Hom}(T\mathcal{E}_z, \mathcal{O}_X(-R))$ over U , and $T\mathcal{E}_z \cong \mathcal{E}_z$ since T is constant. Patching this argument together globally over X gives the isomorphism we desire.

The linearity of L and the tangent equation (16) are proven in [16, Lemma 24 and p 846]. It must satisfy $\mu_*L \cong L^{-1}$ to respect (15) and therefore L takes values in the Prym variety $\text{Prym}(X, \mu) = \{L \in \text{Jac}(X) : \mu_*L \cong L^{-1}\}^0$. $\square$

It is not hard to see that the double periodicity of the map φ obliges L to be doubly periodic. However, it was shown in [18, Thm 5] that L is doubly periodic merely when the induced metric of φ (equally, the solution of (13)) is doubly periodic. As explained in [17, Prop 2], the double periodicity of the map φ is equivalent to the following slightly stronger condition. Let X' be the singular curve obtained from X by identifying the three points O_1, O_2, O_3 lying over $\lambda = 1$. Let $P(X', \mu)$ be the pullback to $\text{Prym}(X, \mu)$ of the group extension $\text{Jac}(X') \rightarrow \text{Jac}(X)$ (whose kernel is isomorphic to $\mathbb{C}^* \times \mathbb{C}^*$) and let

$$P_{\mathbb{R}}(X', \mu) = P(X', \mu) \cap J_{\mathbb{R}}(X')^0; \quad J_{\mathbb{R}}(X') = \{L \in \text{Jac}(X') : \overline{\rho_*L} \cong L^{-1}\}^0.$$

Then φ is doubly periodic iff the real homomorphism

$$(17) \quad L' : \mathbb{R}^2 \rightarrow P_{\mathbb{R}}(X', \mu)$$

is doubly periodic, where this is uniquely specified by the condition that

$$dL'_0\left(\frac{\partial}{\partial z}\right) = d\mathcal{A}'_{P_{\infty}}\left(\frac{\partial}{\partial \zeta^{-1}}\right),$$

with $\mathcal{A}'_{P_{\infty}} : X' \rightarrow \text{Jac}(X')$ the Abel map for X' with base point P_{∞} .

To understand these double periodicity conditions better, let $\Gamma(\Omega')$ denote the space of regular differentials on X' . It is not hard to show that $P_{\mathbb{R}}(X', \mu) \simeq V/\Lambda$ where

$$V = \{\alpha \in \Gamma(\Omega')^* : \rho^*\alpha = -\bar{\alpha}, \mu^*\alpha = -\alpha\},$$

and Λ is the intersection of V with the image of the lattice $H_1(X' \setminus \{O\}, \mathbb{Z})$ in $\Gamma(\Omega')^*$. Here O denotes the singular point of X' over $\lambda = 1$. We can deduce from [19, p 432] that V/Λ is a real compact $(p+2)$ -torus whenever X is smooth. It is easily shown that the image of dL' at the origin is the real two-dimensional subspace $W \subset \Gamma(\Omega')^*$ whose elements annihilate all those regular differentials on X' which vanish at P_0 and P_{∞} . Since Λ is a full lattice, by choosing its generators we may identify V with $\mathbb{R}^{p+2}$. Let us use (W, Λ) to denote the real 2-plane in $\mathbb{R}^{p+2}$ we obtain from this. Then L' is doubly periodic precisely when (W, Λ) is a rational 2-plane, that is, has a basis consisting of rational vectors.

4.2. The moduli of minimal Lagrangian tori. We have learnt that every minimal Lagrangian torus gives rise to spectral data $(X, \lambda, \mathcal{L})$ and we know from [16] that, conversely, any data satisfying the conditions described above gives rise to a minimal torus. It remains to check that the extra symmetry of the holomorphic involution μ on X forces this torus to be Lagrangian as well. This step will complete the proof of the following theorem.

Theorem 1. *There is a bijective correspondence between:*

- (i) *congruence classes of minimal Lagrangian tori $\varphi : T^2 \rightarrow \mathbb{CP}^2$ (with base point on T^2), and;*
- (ii) *equivalence classes of data $(X, \lambda, \mathcal{L}, \mu)$ where, in addition to the reality*

conditions described earlier, $\mathcal{L}$ satisfies (15) and the real homomorphism L' in (17) is doubly periodic.

Proof. This follows from [16, 17, 18] once we have established that (15) forces φ to be Lagrangian. First we must recall how to construct φ from this spectral data. From (X, λ) we construct the real homomorphism $L_z : \mathbb{C} \rightarrow P_{\mathbb{R}}(X, \mu)$ satisfying the tangency condition (16). For each $z \in \mathbb{C}$ we fix the line

$$\ell_0(z) = \Gamma(X, \mathcal{L}_z(-P_0 - P_{\infty})) \in \mathbb{P}\Gamma(X, \mathcal{L}_z) \cong \mathbb{CP}^2$$

where $\mathcal{L}_z = \mathcal{L} \otimes L_z$. The reality condition (14) ensures that $\lambda_* \mathcal{L}_z$ is trivial and therefore $h^0(\mathcal{L}_z) = 3$. To identify all these projective spaces to get a minimal map we need the (essentially unique) trivialising section θ_z of L_z over $X \setminus \{P_0, P_{\infty}\}$ which has the properties: (i) $\theta_z \exp(-z\zeta)$ extends holomorphically to P_{∞} ; (ii) $\theta_z \exp(-\bar{z}\zeta^{-1})$ extends holomorphically to P_0 ; (iii) $\overline{\rho_* \theta_z} = \theta_z^{-1}$, $\mu_* \theta_z = \theta_z^{-1}$. Since $\lambda_* \mathcal{L}_z$ is trivial any global section of $\mathcal{L}_z$ is determined by its restriction to the three points O_1, O_2, O_3 over $\lambda = 1$. So we identify $\Gamma(X, \mathcal{L}_z)$ with $\Gamma(X, \mathcal{L}_0)$ by mapping any global section s of $\mathcal{L}_z$ to the unique global section of $\mathcal{L}_0$ which has the same values as $s\theta_z^{-1}$ at O_1, O_2, O_3 . According to [16, 17] this constructs a minimal map $\varphi : \mathbb{C} \rightarrow \mathbb{CP}^2$ which is doubly periodic precisely when the map (17) is.

To show that φ is Lagrangian it suffices to construct a loop of frames F_{λ} for which F_1 frames φ and α_{λ} possesses the symmetry $\alpha_{-\lambda} = -T\alpha_{\lambda}^t T$. This frame is constructed using the commutative diagram:

$$\begin{array}{ccc} \Gamma(X \setminus \{P_0, P_{\infty}\}, \mathcal{L}_z) & \xrightarrow{\otimes \theta_z^{-1}} & \Gamma(X \setminus \{P_0, P_{\infty}\}, \mathcal{L}_0) \\ f_z \downarrow & & \downarrow f_0 \\ \mathbb{C}(\lambda^{-1}, \lambda) \otimes \mathbb{C}^3 & \xrightarrow{F_{\lambda}^{-1}} & \mathbb{C}(\lambda^{-1}, \lambda) \otimes \mathbb{C}^3. \end{array}$$

The maps f_z are determined by fixing a unitary basis e_j^z for $\Gamma(X, \mathcal{L}_z)$ (the Hermitian metric on this space is described in [16, §3]) for which

$$e_0^z \in \Gamma(X, \mathcal{L}_z(-P_0 - P_{\infty})), \quad e_1^z \in \Gamma(X, \mathcal{L}_z(-2P_{\infty})), \quad e_2^z \in \Gamma(X, \mathcal{L}_z(-2P_0)).$$

The scaling freedom available in the choice of this basis means F_{λ} is only determined up to right multiplication by a diagonal matrix independent of λ . The reality conditions ensure that $F_{\bar{\lambda}^{-1}} = \overline{(F_{\lambda}^{-1})^t}$ over $|\lambda| = 1$ (and therefore everywhere). Now we can use the fact that $\overline{\rho\mu_* \mathcal{L}} \cong \mathcal{L}$ and $\rho\mu(P_0) = P_{\infty}$ to insist that we also have

$$\overline{\rho\mu_* e_0} = e_0, \quad \overline{\rho\mu_* e_1} = e_2,$$

and therefore $F_{-\bar{\lambda}}^{-1} = T\bar{F}_{\lambda}T$. Putting the two symmetries of F_{λ} together shows that $F_{-\lambda} = T(F_{\lambda}^{-1})^t T$ whence $\alpha_{-\lambda} = -T\alpha_{\lambda}^t T$ as required. $\square$

It is now possible to count the dimension of the moduli space of minimal Lagrangian tori in $\mathbb{CP}^2$ (cf. [18, §5], where this is done for minimal tori). At first we will assume that $P_0 + P_{\infty}$ is not a hyperelliptic divisor on X (recall from [18, §5] that this case must be treated differently, for the map φ is S^1 -equivariant then).

First, the data (X, λ, μ) is completely determined by the ramification divisor R . This has degree $2g + 4$ but it must include the divisor $2P_0 + 2P_\infty$. Moreover, it must be preserved by the involutions ρ and μ , neither of which fix any of the points in $R - 2P_0 - 2P_\infty$, so there are $g = 2p$ real parameters available for (X, λ, μ) . The line bundle $\mathcal{L}$ lives in the real p -dimensional space $\{\mathcal{L} \otimes L : L \in P_{\mathbb{R}}(X, \mu)\}$.

Following the discussion at the end of the previous subsection, we can reduce the double periodicity conditions to the study of the map

$$(X, \lambda, \mu) \mapsto (W, \Lambda) \in Gr_{\mathbb{R}}(2, \mathbb{R}^{p+2}).$$

The domain and codomain both have real dimension $2p$. Given local surjectivity of this map (the proof of which has not yet been attempted) we will be able to find spectral data for which the plane (W, Λ) is rational and therefore obtain minimal Lagrangian tori. But where the map is locally surjective it must also be injective. However, for $p > 2$ this does not mean each torus will be isolated, since each torus will live in a real $(p - 2)$ -dimensional family corresponding to the variation of $\mathcal{L}$ (where we have factored out the action of translations along the universal cover of T^2 ; that is, discarded the base point on T^2). Therefore we expect the moduli space to be generically of dimension $p - 2$.

In the case where X is hyperelliptic with hyperelliptic divisor $P_0 + P_\infty$ it follows from [18, §5] that the map φ is S^1 -equivariant. Such minimal Lagrangian tori have been thoroughly discussed [6, 11, 13] and in particular the spectral data approach is unnecessary, since the equation (13) reduces to an o.d.e. which can be solved by elliptic functions. The geometrical explanation for the appearance of elliptic functions is that, from [18, §5], X is generically genus 2 hence $Prym(X, \mu)$ is an elliptic curve.

4.3. The case of genus four spectral curves. In general the construction of examples is hampered by the difficulty in satisfying the periodicity conditions. Once solved, explicit formulae can be found using Riemann θ -functions (see, for example, [21, 19, 15]). When X has genus four the real group $P_{\mathbb{R}}(X, \mu)$ is two dimensional and therefore the corresponding solutions of the Tzitzéica equation are doubly periodic. Indeed, since μ has two fixed points $Prym(X, \mu)$ is principally polarized and therefore the Jacobi variety of a genus two curve. For this reason it is interesting to describe the curves of genus four which can appear.

Proposition 5. *If (X, λ) is the spectral data for a minimal Lagrangian torus in $\mathbb{CP}^2$ and X has genus $g = 4$ then (X, λ) arise from either:*

(i) *the smooth compactification of a curve with equation*

$$(18) \quad \lambda^2 - 2b(x) + x^3\lambda^{-2}, \quad b(x) = b_0 + b_1x + \bar{b}_1x^2 + \bar{b}_0x^3, \quad b_j \in \mathbb{C},$$

(ii) *the smooth compactification of a curve with equation*

$$(19) \quad k\lambda^2 - 2b(x) + k^{-1}\lambda^{-2}, \quad \deg_x b(x) = 3, \quad b(x) \in \mathbb{R}[x], \quad |k| = 1.$$

Proof. From our discussion above we see that the covering $\delta : X \rightarrow Y$ uniquely determines X as the covering curve for $\lambda = \sqrt{y}$ where $y : Y \rightarrow \hat{\mathbb{C}}$ is the rational function of degree three with a single zero at $Q_0 = \delta(P_0)$ and a single pole at

$Q_\infty = \delta(P_\infty)$. All we require of (Y, y) is that it have genus 2 and possess a real involution ρ for which $\overline{\rho^*y} = y^{-1}$ and which fixes every point over $|y| = 1$, on which there are no branch points. Now we recall that every genus 2 curve has a unique hyperelliptic involution: let $\iota \in \text{Aut}(X)$ denote this involution. Since $\rho\iota\rho$ is also a hyperelliptic involution we deduce that $\rho\iota = \iota\rho$. Now we separate the argument into two cases: (i) $\iota(Q_0) \neq Q_\infty$, (ii) $\iota(Q_0) = Q_\infty$.

In case (i) we can choose degree 2 function x with divisor $(x) = Q_0 + \iota(Q_0) - Q_\infty - \iota(Q_\infty)$ and this therefore has $x\overline{\rho^*x}$ constant. We can rescale by a positive constant to make $x\overline{\rho^*x} = 1$. By comparing divisors it follows that $y\iota^*yx^{-3}$ is also constant and by a unimodular scaling of x we may assume this is positive, $y\iota^*yx^{-3} = a > 0$. But

$$a^{-3}x^{-3} = \overline{\rho^*(y\iota^*y)} = a^3x^{-3},$$

so $a = 1$. Thus $y\iota^*y = x^3$. Again by considering divisors we deduce that $y + \iota^*y = 2b(x)$ where $b(x)$ is a polynomial of degree 3. Therefore

$$y - 2b(x) + x^3y^{-1} = 0.$$

From this we compute $\overline{b(\bar{x})} = x^3b(x^{-1})$ so $b(x)$ has the form given in (18).

In case (ii) we choose x to have divisor of poles $Q_0 + Q_\infty$ and so that $\overline{\rho^*x} = x$. In this case $y\iota^*y$ is a unimodular constant k^2 . Set $\tilde{y} = y/k$ then $\tilde{y}\iota^*\tilde{y} = 1$. As above $\tilde{y} + \iota^*\tilde{y} = 2b(x)$ for some polynomial of degree 3. Therefore

$$ky - 2b(x) + k^{-1}y^{-1} = 0, \quad |k| = 1,$$

and a computation shows that $b(x) \in \mathbb{R}[x]$. □

This description does not include the conditions on the ramification divisor R , which are somewhat complicated (however, see the Remark below). In standard hyperelliptic form the corresponding curves Y are as follows. For case (i) set $w = y - b(x)$ then

$$w^2 = b(x)^2 - x^3.$$

For case (ii) set $w = ky - b(x)$, then

$$w^2 = b(x)^2 - 1.$$

It is possible to identify $\text{Prym}(X, \mu)$ with the Jacobian of a genus two curve lying in the symmetric product S^2X , using Donagi's bigonal construction [7]. However, let us finish by simply pointing out that for curves of the type (19) the Prym variety is isogenous to a product of elliptic curves. This echoes the construction of minimal Lagrangian tori by Joyce in [13, §6.2], where he finds formulae which involve products of two different elliptic functions. The spectral curves he produces for these examples are all of type (19).

Proposition 6. *Let E_1, E_2 be the elliptic curves with respective equations*

$$(i) \ w^2 = b(x) + 1; \quad (ii) \ w^2 = b(x) - 1, \quad \deg_x b(x) = 3, \ b(x) \in \mathbb{R}[x].$$

Then for X of type (19), $\text{Prym}(X, \mu)$ is isogenous to $E_1 \times E_2$.

Proof. Any curve X with equation (19) admits two involutions ι_1, ι_2 given by

$$\iota_1(x, \lambda) = (x, \lambda^{-1}); \quad \iota_2(x, -\lambda^{-1}).$$

Notice that $\iota_1 \iota_2 = \mu$. The respective rings of invariants are

$$\mathbb{C}[x, \lambda]^{\iota_1} = \mathbb{C}[x, \lambda + \lambda^{-1}], \quad \mathbb{C}[x, \lambda]^{\iota_2} = \mathbb{C}[x, \lambda - \lambda^{-1}].$$

Taking, respectively, $w = (\lambda + \lambda^{-1})/\sqrt{2}$ and $w = (\lambda - \lambda^{-1})/\sqrt{2}$ yields the equations for E_1 and E_2 . Let $f_j : X \rightarrow E_j$ be the corresponding coverings, then we have the isogeny

$$\text{Jac}(X) \approx f_1^* E_1 \times P_1 \approx f_2^* E_2 \times P_2$$

where P_1, P_2 are the respective Prymians. Since ι_1, ι_2 commute it follows that $f_1^* E_1 \cap P_2$ and $f_2^* E_2 \cap P_1$ are distinct one dimensional subgroups. Now if $L \in f_1^* E_1 \cap P_2$ then

$$\mu^* L = \iota_1^* \iota_2^* L \cong \iota_1^* L^{-1} \cong L^{-1},$$

and similarly for $L \in f_2^* E_2 \cap P_1$. It follows by considering dimensions that

$$\text{Prym}(X, \mu) \approx (f_1^* E_1 \cap P_2) \times (f_2^* E_2 \cap P_1) \approx E_1 \times E_2.$$

□

Remark. There is a simple characterization of the polynomials $b(x) \in \mathbb{R}[x]$ which give spectral data of the type (19); i.e., those polynomials for which the real involution satisfies the right conditions over $|\lambda| = 1$.

Lemma 3. *A pair (X, λ) with equation (19) satisfies all the properties for being spectral data if and only if for some scaling of x*

$$b(x) = x^3 - \frac{3}{2}(u+v)x^2 + 3uvx + w, \quad u, v, w \in \mathbb{R},$$

where $v - u > 4^{1/3}$ and

$$1 + \frac{1}{2}u^2(u - 3v) < w < \frac{1}{2}v^2(v - 3u) - 1.$$

Proof. By rescaling x we may assume $b(x)$ is monic. We must show that these are precisely the conditions under which $\lambda : X \rightarrow \hat{\mathbb{C}}$ has no branch point on $|\lambda| = 1$ and $\rho(x, \lambda) = (\bar{x}, \bar{\lambda}^{-1})$ fixes every point over $|\lambda| = 1$. The locus of points on X over the unit circle is characterized by the equation $b(x) - \cos(\theta) = 0$, $\theta \in [0, 2\pi]$: we see this by setting $\lambda^2 k = e^{i\theta}$ in (19). We require that this equation has distinct real roots for all values of θ . If we write

$$b'(x) = 3(x - u)(x - v), \quad u < v,$$

then since $|\cos(\theta)| < 1$ we require $b(u) > 1$ and $b(v) < -1$: these two give the last inequality in the lemma. In particular $2 < b(u) - b(v) = (v - u)^3/2$, which gives $v - u > 4^{1/3}$. □

4.4. Totally real superconformal maps into $\mathbb{CP}^{2k}$. When n is even those superconformal maps $\varphi : M \rightarrow \mathbb{CP}^n$ which are “totally real” (in the sense that $\varphi^*\omega = 0$, see the remark below) can also be characterized by the existence of a certain primitive harmonic lift. Essentially by definition [2], superconformal maps have an orthogonally periodic harmonic sequence and therefore admit a primitive lift $\psi : M \rightarrow SU_{n+1}/H$, where H is the maximal torus in SU_{n+1} : the target is the full flag manifold over $\mathbb{CP}^n$. Then φ is totally real if and only if it admits a further primitive lift $\tilde{\psi} : \tilde{M} \rightarrow SU_{n+1}/H_0$ where $H_0 \subset H$ is the $n/2$ -dimensional subgroup fixed by the outer involution μ of $\mathfrak{a}_n$ induced by the $\mathbb{Z}_2$ symmetry of its Dynkin diagram. To see this, write $k = n/2$ and order the harmonic sequence $\ell_0, \ell_1, \dots, \ell_n$ so that φ corresponds to ℓ_k . Following [3] we choose local sections f_j of ℓ_j so that

$$\frac{\partial f_j}{\partial z} = f_{j+1} + \frac{\partial}{\partial z} \log |f_j|^2 f_j, \quad j = 0, \dots, n-1.$$

These imply the Toda equations

$$\frac{\partial^2}{\partial z \partial \bar{z}} \log(|f_j|^2) = \frac{|f_{j+1}|^2}{|f_j|^2} - \frac{|f_j|^2}{|f_{j-1}|^2}, \quad j = 0, \dots, n-1.$$

Since φ is totally real we can arrange for $|f_k| = 1$. An inductive argument using the Toda equations then shows that $|f_{k+j}| = |f_{k-j}|^{-1}$ for $j = 0, \dots, k$. Now let $F : \tilde{M} \rightarrow SU_{n+1}$ be the Toda frame, whose j -th column is $f_j/|f_j|$, and let $\alpha = F^{-1}dF$ be its Maurer-Cartan form. Then a simple computation shows that $\alpha_{\mathfrak{h}}$ lies in $\mathfrak{g}_0^\nu \cap \mathfrak{g}_0^\mu$ and $\alpha_{\mathfrak{p}}^{1,0}$ lies in $\mathfrak{g}_1^\nu \cap \mathfrak{g}_1^\mu$, where ν is again the Coxeter-Killing automorphism. Therefore, with an appropriate initial condition for F , all the maps in this harmonic sequence have a (local) primitive lift into SU_{n+1}/H_0 , since this is the $2n+2$ -symmetric space for the product automorphism $\nu\mu$.

In particular, for totally real superconformal tori we can argue as above that the lift factors through a covering torus, with covering group a subgroup of $\mathbb{Z}_{n+1}$. The spectral data for such tori will consist of $(X, \lambda, \mathcal{L}, \mu)$ where μ is a holomorphic involution which the other data respects in the same fashion as for the case $n = 2$.

Remark. There is a schism in the mathematical community as to the definition of a totally real submanifold. It seems that the original meaning, used in the 1960’s in the study of real submanifolds M of a complex submanifold, is that $TM \cap J(TM) = 0$, where J is the complex structure on the ambient space. Somewhere in the 1970’s this term started to be used also for the condition that TM is perpendicular to $J(TM)$, which is the sense I have used above. Unfortunately, we now have the situation where both definitions are in use and it is not always clearly stated which definition is meant.

REFERENCES

- [1] M Black, *Harmonic maps into homogeneous spaces*, Pitman Research Notes in Math. 255, Longman, Harlow 1991.
- [2] J Bolton, F Pedit & L Woodward, *Minimal surfaces and the affine Toda field model*. J. reine angew. Math. 459 (1995), 119-150.
- [3] J Bolton & L Woodward, *Congruence theorems for harmonic maps from a Riemann surface into $\mathbb{CP}^n$ and S^n* , J. London Math. Soc. 45 (1992), 363-376.

- [4] F Burstall, *Harmonic tori in spheres and complex projective spaces*, J. reine angew. Math. 469 (1995), 149-177.
- [5] F E Burstall & F Pedit, *Harmonic maps via Adler-Kostant-Symes theory*, in *Harmonic maps and integrable systems*, ed: A P Fordy & J C Wood, Aspects of Mathematics E23, Vieweg 1994.
- [6] I Castro & F Urbano, *New examples of minimal Lagrangian tori in the complex projective plane*, Manuscripta Math. 85 (1994), 265-281.
- [7] R Donagi, *The fibers of the Prym map*, in *Curves, Jacobians and abelian varieties*, Contemp. Math. 136, 55-125, AMS 1992.
- [8] J Eells & L Lemaire, *Selected topics in harmonic maps*, CBMS Regional Conference Series in Mathematics 50, AMS 1980.
- [9] D Ferus, F Pedit, U Pinkall & I Sterling, *Minimal tori in S^4* , J. reine angew. Math. 429 (1992), 1-47.
- [10] M Gross, *Special Lagrangian fibrations II. Geometry. A survey of techniques in the study of special Lagrangian fibrations*, Surv. Diff. Geom. 5 (1999), 341-403.
- [11] M Haskins, *Special Lagrangian cones*, arXiv:math.DG/0005164 (2000).
- [12] R Harvey & H B Lawson, *Calibrated geometries*, Acta Math. 148 (1982), 47-157.
- [13] D Joyce, *Special Lagrangian 3-folds and integrable systems*, to appear in Proceedings of the 9th Math. Soc. Japan Internat. Res. Inst. arXiv:math.DG/0101249.
- [14] D Joyce, *Singularities of special Lagrangian fibrations and the SYZ conjecture*, arXiv:math.DG/0011179 (2000).
- [15] Hui Ma & Yujie Ma, *Totally real minimal tori in $\mathbb{CP}^2$* , arXiv:math.DG/0106141 (2001).
- [16] I McIntosh, *The construction of all non-isotropic harmonic tori in complex projective space*, Internat. J. Math. 6 (1995), 831-879.
- [17] I McIntosh, *Two remarks on the construction of harmonic tori in $\mathbb{CP}^n$* , Internat. J. Math. 7 (1996), 515-520.
- [18] I McIntosh, *On the existence of superconformal 2-tori and doubly periodic affine Toda fields*, J. Geometry Phys. 24 (1998), 223-243.
- [19] I McIntosh, *Harmonic tori and generalised Jacobi varieties*, Comm. Anal. Geom. 9 (2001), 423-449.
- [20] B O'Neill, *The fundamental equations of a submersion*, Michigan Math. J. 13 (1966), 459-469.
- [21] R Sharipov, *Minimal tori in the five dimensional sphere in $\mathbb{C}^3$* , Theoret. and Math. Phys. 87 (1991), 363-369.
- [22] A Strominger, S-T Yau & E Zaslow, *Mirror symmetry is T-duality*, Nuclear Phys. B 479 (1996), 243-259.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK YO10 5DD, UK

E-mail address: im7@york.ac.uk